

**Publication:** **Фармакология - Практическому здравоохранению, Том 7, Сентябрь 2007, Санкт-Петербург**  
*Pharmacology – Practical Healthcare, Issue No: 7, September 2007*  
*St. Petersburg, Russia*

**Authors:** **Луспикаян С. Х., Тараканов А. В., Воронкин Д. А.**  
**(Luspikeyan S.H., Tarakanov A.V., Voronkin D.A.)**  
*Ростовский государственный медицинский университет*  
*Росздрава, Нахичеванский пер.,29, Ростов-на-Дону, 344022*  
*State medical university of Rostov, 29, Nahichevan st., Rostov-on-Don, 344022*

**Article name:** **Многофакторный механизм СКЭНАР-Анальгезии**

## **Effect of arthrophon with use of Transcutaneous Neurostimulation on values of lipid peroxidation at complex post-surgical therapy in patients with purulent appendicular peritonitis**

The tumour necrosis factor-alpha (TNF) plays a significant part in inflammatory destruction of pathogenesis.

Purpose – to explore the influence of arthrophon in respect of its bipathic effect combined with transcutaneous neurostimulation (SCENAR-therapy) on the state of lipid peroxidation, values of endogenous intoxication at treatment of patients with purulent appendicular peritonitis in post-surgical period.

Materials and methods: 58 patients were examined who were operated for purulent peritonitis. All the patients were treated with conventional intensive therapy. In group 1 (n =38) treatment combined with SCENAR was performed. In group 2 (n=20), four tablets per day of arthrophon were additionally prescribed, plus 5 procedures of SCENAR-therapy. The control (n=36) – healthy donors comparable as age and sex. Evaluation of efficiency was performed according to clinical and laboratory data on the first and fifth day from the start of the treatment. Application of arthrophon led to decrease of the level of myeloperoxidase with 50.1% ( $p<0.001$ ). The activity of catalase increased with 41% compared with the initial background ( $p<0.02$ ) and reached the control group. In both groups, statistically reliable decrease of malondialdehyde (MDA) was noted as well as decrease of diene-conjugates on the 5th day of the treatment. The level of intoxication was defined according to the fraction of medium-mass molecules in blood plasma. After a five-day course of therapy in the first group the contents of medium-mass molecules (280) decreased with 28.8% ( $p<0.01$ ), in the second group – with 39% ( $p<0.001$ ). In the second group, decrease of medium-mass molecules was noted – 30.3% ( $p<0.05$ ), in the first group a reliable change of this value was not noted. On the background of arthrophon and SCENAR-therapy, the level of circulating immune complex dramatically decreases 5 days after the treatment – 40.3% ( $p<0.01$ ).

Conclusions: arthrophon combined with SCENAR-therapy proves to be an effective therapeutic means for post-surgical period in patients with purulent appendicular peritonitis. A significant decrease of clinical symptoms of the disease was noted, as well as normalization of paraclinical values. The combined therapy allows to correct faster and more efficiently the extent of endogenous intoxication, which were evaluated with reliable decrease of some fractions of medium-mass molecules and circulating immune complex. Arthrophon and SCENAR-therapy combined with basic intensive therapy in post-surgical period in patients with purulent appendicular peritonitis decrease the intensity of processes of lipid peroxidation on the 5th day, increase anti-oxidant protection and improve the function condition of cellular membranes.

### Summary:

**The complex effect of SCENAR-therapy is demonstrated in situations of emergency, when various conditions associated by acute pain are treated. The result of monotherapy of pains: proved analgesic effect at traumas (n=20, 60%); analgesia against a background of normalized function of the system of organs at stenocardia (n=26, 66%) and hypertensive crises (n=153, 88-91% for cephalgia, and cardialgia respectively).**

The issue of protecting patients against pains has an enormous humanistic importance. Pain is a signal of danger, a symptom of many diseases, main reason for patients to suffer, basic pathophysiological factor of developing shock.

When a patient faces aversive factors of ambience, adaptive mechanisms start so that the organism survives in new conditions. Activation of endogenous antinociception system is one of these general biologic mechanisms. It is proved that realization of endogenous antinociception is performed through opiate and non-opiate systems of brain. Mechanisms of antinociception have been subject of numerous comprehensive works, including acupuncture, electropuncture, and transcutaneous neurostimulation. [2,3,8,11]

It is known that pharmaceutical methods of analgesia are imitation of the functions of endogenous system to protect against pain. They are important when performing mass analgesia. Starting the endogenous antinociception system is optimal when methods of reflexotherapy are used, and this method is more physiologic. The complexity of the methods, their invasiveness, impossibility to be used in extreme circumstances, lack of reliable and inexpensive equipment, and insufficiency of trained specialists are the factors that obstruct mass introduction of reflexological methods of analgesia.

Methods of transcutaneous bioregulated low-frequency impulsive electrotherapy have been used recently for non-pharmaceutical treatment [5,6,7], particularly – self-controlled energy neuroadaptive regulator (SCENAR). This method is based on the effect of bipolar impulsive points of low frequency without constant component. Characteristics which distinguish SCENAR among electric equipment for transcutaneous stimulation: 1) optimal form of the impulse and the force of the effect; 2) the body almost does not have to adapt to SCENAR impact due to biotechnological feed-back; 3) non-damaging mode of impact; 4) high level abruptness of the front of the affecting signal with neutralization of the accommodation effect; 5) many researches have been performed to examine this method of treatment; 6) during the treatment an expert

evaluation of the dermal condition can be used to optimize the area of impact; 7) optimal construction of the appliance enables it to work in various conditions [6, 7].

Applying SCENAR-therapy is an issue of huge interest due to its availability, rapid positive effect, simplicity of treatment and steady result.

## **TASKS OF THE RESEARCH**

Examining possibilities to perform SCENAR-therapy at various pathological processes accompanied by acute pain in conditions of emergency.

## **MATERIALS AND METHODS**

Mobilizing adaptive systems of the organism using SCENAR in conditions of acute damage of their functions and analgesic effect were the reason to use the appliance in emergency situations. More than 40 doctors of emergency departments in 13 towns of district of Rostov, Stavropol, Nizhni Novgorod and Adigeia took part in this research. More than 500 patients were helped. The coordination of clinical tests according to special developed map was performed by Emergency department of Faculty of qualification improvement of State Medical University of Rostov and the department of clinical tests of "RITM" Development Bureau, town of Taganrog. Patients with acute pain at three different pathological processes were selected from the database: 1) traumatic injuries of limbs, soft tissue injuries (n=20); 2) steady stenocardia (n=26); 3) hyper-intensive crises (n=153).

SCENAR-therapy was performed according to instructions [12], as dermal areas with located "anti-pain" points were included; if needed methods of general treatment can be added: "three paths on the backbone and six points on the face" and "collar area, forehead, adrenals". If there was no possibility to affect the injured organ, "work" with the appliance was performed on symmetric areas. The kin was treated using CHENS-SCENAR appliances without digital methods in mode F1. The therapeutic procedure was performed only once by an emergency department doctor, as the duration of the procedure was 5-30 minutes long, depending on the effect. When the analgesia was ineffective, the doctor could inject analgesic medicine. To evaluate the effect of the analgesia in conditions of emergency, a visual analog scale (VAS) was used, where 0cm = lack of pain, 10cm = maximum pain.

The use of itineraries subject to the method "three paths" of the backbone and six points of the face was performed on the grounds of the following: when processing path 2 and 3, Shu-points of the back are included in the area of the impact of SCENAR (points of consent)[4, 10, 11], which are located on the urethra. The itinerary of path 1 is a non-pair rear middle meridian. When areas containing points VG1-VG4 are processed a therapeutic effect is marked on this channel at nervous system diseases, intestinal diseases, diseases of the urino-genital system. When areas containing points VG5-VG8 are processed a therapeutic effect is marked on this channel at nervous system diseases, diseases of gastric and intestinal system. The area of points from VG9 to VG14 - nervous system disorders, pulmonary diseases, fever.

Processing six points on the face enables to input information through all the three ramifications of the trigeminus nerve. On the other hand biologically active points of the channels are located in these areas. At least point V2 of the urethra, points E2 and E3 of the gastric channel and G120 of the large intestine, and point E4 of the gastric channel are included in the area of impact of SCENAR.

The method "collar, forehead, adrenals" is also related to the areas of general treatment. The use of itineraries in this method is performed due to largely represented reflexogenic areas and biologically active points as the experience of SCENAR-therapy shows [5, 12]. When the indicated itineraries are processed and when the dermal areas are processed "according to instructions", the appliance may stick. In this case the electrode is not unstuck from the skin but is held on it until further movement becomes possible. If the appliance does not stick, then other kinds of so-called asymmetry can appear (hyperimia, change of the characteristics of the sound, different sensibility etc.). These dermal areas are processed additionally until initial characteristics are changed.

## RESULTS AND DISCUSSION

**1. Pain at traumas.** Analgesia with the use of SCENAR was performed at small and medium traumas: lower limbs fracture (n=5), injuries of soft tissues – limbs, thorax, without damaging respiratory functions (n=9), I and II degree burns of face and abdominal wall (n=2). Cut wounds and bites (n=4). There were nine men and 11 women with age from 11 to 82 (average age –  $40.7 \pm 3.9$ ). The patients were treated using general means of treatment: splinting, immobilization, non-adhesive bandages in cases of burn etc. The evaluation of the pain was performed using VAS before the analgesia with SCENAR, immediately after the procedure, and in the 10th and 20th minute. In the same periods of time other complaints and symptoms were defined, as well as the parameters of hemodynamics. The data is shown on fig.1.




Fig.1. **A** - Dynamics of the analgesic effect of SCENAR during therapy of traumatic pains according to VAS.

**B** – Dynamics of systolic and diastolic blood pressure and pulse frequency of patients with traumatic pains. Reliability referring to data before SCENAR: \* -  $P < 0.05$ ; \*\*\*-  $P < 0.001$ ;

A distinct reliable analgesic effect developed immediately after the procedure, almost without further increase by the 20<sup>th</sup> minute. Observations showed that the analgesic effect increases by the 50<sup>th</sup> – 60<sup>th</sup> minute. Because the emergency staff cannot follow the efficiency of the procedure for a long period of time, in several cases the doctor made the decision to inject additionally non-opiate analgesic medicine (analgin, ketorolac or ketonal). Only 8 out of 20 patients were injected analgesic medicines. Natural skepticism and caution should be noticed when doctors tested the new method.

We regard as an important factor the steadiness of the values of blood pressure. Reliable decrease of the average frequency of cardiac contractions from 85 to 81 beats per minute indirectly shows sympathetic tonus. Reliable change of respiratory

frequency was not noticed and it was within normal limits. Decrease of other symptoms and complaints was noticed along with the analgesic effect, such as dizziness, nausea, excitedness and sensation of inhibition.

Independent analgesic effect developing on the background of steady parameters of blood pressure and decreasing of symptoms accompanying the traumas is noticed during the process of analgesia using SCENAR for patients with small and medium traumas.

## 2. Pain at steady stenocardia tension of I-III functional class.

26 patients with ischemic cardiac disease, crises of pain at steady tension stenocardia of different functional classes were treated with SCENAR, as the patients had previous idiosyncrasy to nitrates or lack of efficiency of the nitrates. There were 14 men (average age –  $63,1,7 \pm 2,5$  years old), 12 women (average age -  $68.4 \pm 3.7$  years old). 7 patients had arterial hypertonia as accompanying disease, and 3 of them had osteochondrosis of the backbone. This category of patients was additionally treated with SCENAR as the dermal area of the pericardial channel was processed in the lower part of right forearm and the painful area of skin on the thorax.





Fig.2. **A** – Dynamics of the complete analgesic effect and number of residual pains (%) during analgesia of patients with attacks of stenocardia.  
**B** - Dynamics of analgesia of patients with residual pains during SCENAR-therapy, according to VAS. Reliability referring to SCENAR data: \* -  $P < 0.05$ ; \*\* -  $P < 0.02$ ; \*\*\* -  $P < 0.001$ .

As can be seen in fig.2 (A) immediately after SCENAR-treatment the pain completely disappeared for 46 % of the patients and for 65% of the patients – 20 minutes later. 35% of the patients felt residual pain in the 20<sup>th</sup> minute of the treatment, but the intensity of the pain was much less, which can be seen in fig 2 (B). These patients, among whom there were mainly patients with accompanying diseases, were offered additional therapy with included symptomatic medicines and non-opiate analgesics. Changes of parameters of blood pressure and pulse statistically did not differ for the category of patients with additional treatment and patients with SCENAR-therapy.

This way, analgesia using SCENAR for algic crises at steady tension stenocardia causes sufficient independent analgesic effect, which does not depend on parameters of general pressure. The extent of its correlation with coronary varicose and the endogenous mechanisms which SCENAR starts in order to stop the pain, can be defined in hospital conditions and additional tests.

### **3. Cardialgia and cephalgia at hypertensive crises.**

During SCENAR-therapy for patients with neurovegetative form of hypertensive crises normalization of parameters of blood pressure was aimed, not only analgesic effect. A total of 153 patients were helped at their homes on emergency call. To analyze the analgesic effect we divided the patients into 2 groups: with cephalgia (n=137, which forms 90% of the patients; among them 71 patients – 52% had cardialgia); separately data of patients with cardialgia were analyzed (n=78, which is 51% of the patients). Patients of this category were treated by SCENAR, as the skin was processed in the collar area, area of pericardial channel, in the lower part of the right forearm, and dermal projections of maximal pain.



**Fig.3. A** – Dynamics of the complete analgesic effect and number of residual pains (%) during analgesia of patients with cardialgia at hypertensive crisis.


**B** - Dynamics of analgesia of patients with residual pains during SCENAR-therapy, according to VAS. Reliability referring to SCENAR data: \* -  $P < 0.05$ ; \*\* -  $P < 0.02$ ; \*\*\* -  $P < 0.001$ .

In fig. 3 can be seen that immediately after the procedure cardiac pains disappeared for 61% of the patients, and for 91% by the 30<sup>th</sup> minute. Nevertheless, residual painful sensations were registered by the 30<sup>th</sup> minute only for 9% of the patients. Their extent was much lower, as at stenocardia, which could be seen during evaluation using visual analog scale (fig. 3 B). This category of patients at times needed an additional injection of non-opiate analgesic.

In fig.4, the effect of the appliance can be seen for cephalgia and hypertensive crises at pre-hospital stage of treatment. Fig.4 A shows reliable and gradual elimination of pain. After the procedure the headache disappeared for 43%, and for 88% of patients - by the 30<sup>th</sup> minute.

Residual pains by the 30<sup>th</sup> minute were registered only for 12% of the patients. Their extent, according to VAS, was much lower than before the procedure (fig.4 B). Unidirectional elimination of pain in cardiac area and headaches was noticed during elimination of hypertensive crisis. We analyzed the changes of parameters of blood pressure during SCENAR-therapy.

Some patients took hypotensive medicines before the emergency staff arrived, which dramatically changes the efficiency of SCENAR. Naturally there was certain skepticism: was it SCENAR that affected or was it previously taken medicines?




**Fig.4. A** – Dynamics of the complete analgesic effect and number of residual pains (%) during analgesia of patients with cephalgia at hypertensive crisis.

**B** - Dynamics of analgesia of patients with residual pains during SCENAR-therapy, according to VAS. Reliability referring to SCENAR data: \* - P < 0.05; \*\* - P < 0.02; \*\*\* - P < 0.001.


This is why we divided all the patients into 2 groups: ones who took medicines and ones who did not take medicines before they called the emergency and were treated with SCENAR. Among the patients who had taken hypotensive medicines before the emergency arrived (calcium antagonists,  $\beta$ -adrenoblockers, APF inhibitors etc) 30-90 minutes before that without any effect there were 80 people (58 women, 22 men; average age –  $61.1 \pm 4.9$  years old). The group of patients who did not take medicines before calling the emergency contained 73 people (52 women, 21 men; average age –  $63.1 \pm 3.5$  years old). The results are shown in fig.5.

As can be seen in the left part of the figure, the parameters of blood pressure for both groups did not differ before and after SCENAR-treatment, on the background of general decrease of blood pressure. 30 minutes before the end of the procedure the systolic blood pressure decreased 11-16% , diastolic – 11-9%. This speed of decrease of blood pressure in the first 30 minutes is effective as cardialgia and cephalgia are eliminated it is safe for all ages. A typical effect of the SCENAR-therapy is the primary elimination of the complaint of the symptom and then normalization of objective parameters. Vegetative corrective effect of SCENAR-therapy should be noticed as hyperhydrosis, nausea, vomiting and excitedness is eliminated. Further observation of some patients and the experience of elimination crises in hospital conditions show that average blood pressure decreases 15-25% within 1 hour, which does not lead to ischemia of targeted organs. The reliable decrease of frequency of cardiac contractions with 9% could be noticed when eliminating crises without previous use of medicines (fig.5b)

**A**



**B**



**Fig.5. A** – Dynamics of parameters of blood pressure during SCENAR-therapy of hypertensive crises in the group of patients who had taken and who had not taken hypotensive medicines before the emergency staff arrived.  
**B** - Dynamics of frequency of cardiac contractions in the same groups. Reliability referring to the data before SCENAR: \* -  $P < 0.05$




Fig.6. Dynamics of correlation coefficient depending on the decrease of systolic blood pressure and decrease of painful sensibility at cephalgia and cardialgia, according to VAS.

If during the process of analgesia the analgesic effect develops independently for patients with traumas, then during attack of pain at steady stenocardia and particularly hypertensive crises, the analgesia probably depends mostly on recovering of functions and decrease of ischemia: the decrease of system blood circulation and normalization of regional blood circulation including the coronary one.

We analyzed the dynamics of correlation coefficient depending on the decrease of systolic blood pressure and the extent of headache and cardiac ache, according to visual analog scale. The data is represented in fig.6. As can be seen in this figure the correlation coefficient increases simultaneously with the decrease of the systolic blood pressure and decrease of values of extent of cardialgia and cephalgia, according to VAS. If in the beginning of the crisis the extent of the pain did not depend on the values of blood pressure, then SCENAR-therapy led to increase of the correlation to 0.41 and 0.48. Similar dynamics of close relation and its lack in the beginning of the crisis probably shows simultaneously developing effects of SCENAR: analgesic effect and the effect, which normalizes the functional system of blood pressure regulation. This way, the elimination of neurovegetative form of hypertensive crises using SCENAR regardless of preliminary therapy leads to gradual decrease of parameters of blood pressure and frequency of cardiac contractions. Distinct analgesic effect of cephalgia and cardialgia is noticed, which probably has a mixed nature.

During the process of elimination of acute pain with different origins, three elements should be present in the algorithm: possible elimination of reason causing pain; affecting the sensation of pain - perception; possible prophylactics of pathophysiological damages, which pain causes or will cause.

As the experience of work with SCENAR shows, during the treatment of acute and chronic painful syndromes, the data presented in this article show the multicomponentness of the suggested therapy. Non-medical method of elimination of acute pain starts the endogenous antinociception and other mechanisms of recovering damaged functional systems. The analysis of the data obtained in traumatology and cardiology shows that the share of perceptual component of SCENAR analgesia is

different depending on nosology, the level and the volume of the damage. It is not reasonable to rely on only one single method of analgesia and it is also dangerous for the patient, especially in case of emergency.

Undoubtedly, the morphologic substrate of the analgesic effect of SCENAR-therapy are the antinociceptional systems of brain, started from the different areas of skin. SCENAR works with the biologically active points, with their accumulation and the surrounding tissues – this a zonal appliance. Due to its technical solution, it “finds” these points independently. As shown for acupuncture analgesia [8], in opioidergic, serotonergic, catecholaminergic, cholinergic, gamkergetic and possibly other mechanisms take part in its realization.

Some authors regard the transcutaneous electro-stimulation as the most accessible and the simplest “hyperstimulation analgesia” to be realized. If we proceed from the theory of Melzack and Wall (1965) the activity in the thin non-myelinated C-fibers located in the jelly substance of the rear horns of the spinal cord “opens the gates” to the further pass of nervous impulses bearing the information of pain. The role of the thick fibers consists in the ability to “close the gates”. One of the hypotheses of the analgesic effect of transcutaneous neurostimulation consists of the fact that the procedure leads to activation of thick myelinic fibers with subsequent “locking the painful entrance”. The analgesic effect during the so-called dynamic neurostimulation is proved to be naloxone-dependent [9].

Using almost the same methods of SCENAR-therapy and taking under consideration the principles of heir application [5,6,12], we obtained unidirectional results: analgesic effect and final positive sanogenic result connected with normalization of the changed functional system. Taking under consideration the limited volume of such a publication we can mention that similar unidirectional results were obtained during the process of therapy of acute myocardial infarction, burns, bronchial asthma of adults and infants etc.

The simplicity of use of SCENAR allows us to recommend the use of this kind of analgesia in any conditions and if needed, by people without medical education.

## CONCLUSIONS

1. The SCENAR-therapy (self-controlled, energy neuro-adaptive regulator) is an effective, safe and multifactor non-pharmaceutical method of analgesia during treatment of different conditions accompanied by acute pain in cases of emergency.
2. During the process of analgesia using SCENAR the patients with small and medium trauma (n=20), an independent analgesic effect is noticed (60%) developing on the background of steady parameters of blood pressure and decrease the symptoms accompanying traumas.
3. The analgesia of algic attacks at ischemic disease and steady tension stenocardia (n=26) using SCENAR, causes a sufficient independent analgesic effect (66%), which does not depend on the parameters of general pressure.
4. The elimination of neurovegetative form of hypertensive crises (n=153) using SCENAR, regardless of preliminary medical therapy leads to gradual reliable decrease of the frequency of cardiac contractions and decrease the parameters of blood pressure by the 20th minute after the procedure; by the 30<sup>th</sup> minute a distinct analgesia of cephalgia and cardialgia is noticed (88-91% respectively).

## Bibliography:

1. Абабян Г.Н. Рефлексотерапия заболеваний нервной системы. М., 1999. - 297с.
2. Барашков Г.Н. Рефлексотерапия боли. – М.:НЦМ "Универсимед", изд-во "ТМ – ОКО", 1995. – 263с.
3. Василенко А.М. Акупунктура и рефлексотерапия:эволюция методологии и метатеории. – Таганрог. 1998. – 110с.
4. Вогралик В.Г., Вогалик М.В. Акупунктура. Основы традиционной восточной рефлексодиагностики и пунктурной адаптационно-энергизирующей терапии: Цигун. – М.:ГОУ ВУНМЦ РФ, 2001. – 336с.
5. Горфинкель Ю.В. – Теоретические и практические основы повышения эффективности СКЭНАР-терапии. В сборнике статей "СКЭНАР-терапия и СКЭНАР-экспертиза", выпуск 2, Таганрог, 1999. – С. 16-18
6. Гринберг Я.З. К вопросу об основании эффективности СКЭНАР-терапии/СКЭНАР – терапия и СКЭНАР экспертиза. – Таганрог 1997. – Вып.3. – с.16-22.
7. Гринберг Я.З. Концепция электротерапии / СКЭНАР-терапия и СКЭНАР-экспертиза. – Таганрог, 1999. – вып.5 – с.16-13
8. Игнатов Ю.Д., Качан А.Т., Васильев Ю.Н. Акупунктурная аналгезия:Экспериментально-клинические аспекты. Л.Медицина, 1990. – 256с.
9. Кукушкин М.Л., Мейзеров Е.Е., Графова В.Н. и др. Особенности развития анальгетического эффекта при чрезкожной динамической электроннойростимуляции // Бюллетень экспериментальной биологии и медицины. – 2003. - №3. – С.265-267
10. Лувсан Гаава. Очерки метода восточной рефлексотерапии. Новосибирск: Наука, 1991. – 432с.
11. Мачарет Е.Л., Коркушко А.О. Основы электро- и акупунктуры – Киев:Здоров'я, 1993. – 391с.
12. Ревенко А.Н., Тараканов А.В. и др. Инструкция по применению чрезкожного электроннойростимулятора с биообратной связью индивидуально-дозированного воздействия на рефлекторные зоны. ЧЭНС-СКЭНАР. – Таганрог, 2004. 70с.

**The article is provided with the assistance of:**

RITM OKB ZAO  
Petrovskaya 99,  
Taganrog 347900  
Russian Federation  
[www.scenar.com.ru](http://www.scenar.com.ru)

and

RITM Australia Pty Ltd  
4/130-134 Pacific Hwy  
Greenwich NSW 2065  
Australia  
[www.scenar.com.au](http://www.scenar.com.au)

Translated from Russian to English Language by: Mrs. Veselina Petkovap August 2008, Mladost 329, Sofia 1712, Bulgariap <http://syntagma.dir.bg>